

Exact Microscopic Entropy of Non-Supersymmetric Extremal Black Rings

Roberto Emparan

*Institució Catalana de Recerca i Estudis Avançats (ICREA),
and*

*Departament de Física Fonamental, Universitat de Barcelona,
Martí i Franquès 1, E-08028 Barcelona, Spain*

`emparan@ub.edu`

Abstract

In this brief note we show that the horizon entropy of the largest known class of non-supersymmetric extremal black rings, with up to six parameters, is exactly reproduced for all values of the ring radius using the same conformal field theory of the four-charge four-dimensional black hole. A particularly simple case is a dipole black ring without any conserved charges. The mass gets renormalized, but the first corrections it receives can be easily understood as an interaction potential energy. Finally, we stress that even if the entropy is correctly reproduced, this only implies that one sector of chiral excitations has been identified, but an understanding of excitations in the other sector is still required in order to capture the black ring dynamics.

Motivation. Non-supersymmetric extremal rotating black holes provide a convenient setting for furthering our microscopic understanding of quantum processes related to Hawking radiation. On the one hand, they exhibit the attractor behavior that allows precise control over the statistical counting of the entropy. On the other hand, they typically possess ergoregions and thus emit quantum radiation in a process closely related to Hawking radiation, even if they are at zero temperature [1]: this is the spontaneous superradiant emission recently studied in a microscopic model in [2]. Thus, these black holes allow rather detailed control over their microscopic description, while exhibiting richer and more realistic dynamics than supersymmetric black holes.

The non-supersymmetric extremal black *rings* [3, 4, 5] that we discuss here present further advantages in these respects over other non-BPS extremal black holes. The ratio R_2/R_1 between the S^2 and S^1 radii of the ring controls the departure from the BPS limit, which is recovered as $R_1 \rightarrow \infty$. Even if the mass is renormalized as a function of this parameter, we shall argue that in fact such a renormalization can be understood in simple terms in an expansion in R_2/R_1 . In contrast, the entropy is exactly reproduced for *all* values of R_2/R_1 . This result matches nicely with the attractor behavior discussed in [6, 7] and provides a non-trivial test of it. Very recently, the exact entropy of a vacuum black ring, with horizon topology $S^1 \times S^2$ and maximal rotation on the S^2 [8], has been reproduced following on ideas in [9, 10, 11]. However, in this case the connection between the dual CFT and the black hole degrees of freedom is obscured by the series of transformations involved in the mapping. In our examples, the connection to the CFT is much more direct. Still, they present a number of novel features, yet to be fully understood, largely due to the fact that, in contrast to supersymmetric black holes and black rings, these black rings can emit quantum radiation.

Thus we believe that understanding in detail the microscopic dual of these black rings can significantly enlarge the range of black hole phenomena we can address using string theory. As a first step in the correct identification of this dual description, in this note we show how their entropy is exactly reproduced by a microstate counting.

Extremal dipole black rings. Our black rings appear at the common intersection of three stacks of M5 branes over a circle. We begin with the simplest example of these, constructed in [3], which does not possess any conserved charges, only three dipoles q_i associated to each kind of M5 brane. Thus the solutions can never be supersymmetric, but they admit an extremal (zero-temperature) limit with a regular horizon of finite area. Ref. [3] proposed to account for their entropy using the MSW conformal field theory [12], and indeed found perfect agreement for the entropy up to the next-to-leading order in an expansion in R_2/R_1 . Although this is already a non-trivial matching, ref. [3] failed to realize that the matching is exact for arbitrary values of R_2/R_1 , as we shall presently see. Later, the idea to apply the MSW CFT to black rings was shown to work for supersymmetric black rings [13, 14, 15]. However, these calculations left a number of puzzling issues; we will discuss some of them after we deal with the more general solutions of [4], which have M2 charges.

The extremal dipole black rings with three M5 dipoles are recovered from the solutions in [3] by setting the parameter ν to zero. The solutions are expressed in terms of four dimensionless parameters $\lambda, \mu_{1,2,3}$, with $0 < \lambda, \mu_i < 1$. Mechanical equilibrium of the ring then determines that

$$\lambda = \frac{\mu_1 + \mu_2 + \mu_3 + \mu_1\mu_2\mu_3}{1 + \mu_1\mu_2 + \mu_1\mu_3 + \mu_2\mu_3}. \quad (1)$$

The solutions also contain a scale R . Roughly, R measures the size of the ring's S^1 radius, $R \sim R_1$; λ is roughly the ratio between the S^2 and S^1 radii, *i.e.*, $R_2 \sim \lambda R$. Finally, the μ_i are ratios between ‘dipole-charge radii’ and R . A straight string is recovered as $\lambda, \mu_i \rightarrow 0$ keeping λR and $\mu_i R$ finite.

The Bekenstein-Hawking horizon entropy is

$$S_{BH} = \left(\frac{2\pi R^3}{G} \right) \pi \sqrt{\lambda(1-\lambda^2)} \prod_{l=1}^3 \mu_l^{1/2} (1 + \mu_l), \quad (2)$$

and the integer-quantized M5 dipoles are

$$n_i = \epsilon_i \left(\frac{2\pi R^3}{G} \right)^{1/3} \mu_i \frac{\sqrt{1-\lambda}}{\sqrt{1-\mu_i^2}} \prod_{l=1}^3 \sqrt{1+\mu_l}. \quad (3)$$

The rings also possess an angular momentum along the S^1 direction

$$J_1 = \epsilon_\lambda \frac{1}{4} \left(\frac{2\pi R^3}{G} \right) \lambda \sqrt{1+\lambda} \prod_{l=1}^3 (1+\mu_l)^{3/2}. \quad (4)$$

We have included a choice of sign $\epsilon_\lambda, \epsilon_i = \pm 1$ for later generality. For our purposes at finite R_2/R_1 (*i.e.*, finite λ, μ_i) the orientations of the momentum and the M5 branes may remain arbitrary. It is only if we want to recover a supersymmetric solution in the straight-string limit that we must correctly align the momentum and branes.

A direct calculation shows that, for *all* allowed values of the parameters, the entropy (2) can be written as

$$S_{BH} = 2\pi \sqrt{|J_1 n_1 n_2 n_3|}. \quad (5)$$

One naturally expects that if we view the ring as a circular string, then J_1 must be identified as the units n_p of linear momentum running along the string—detailed arguments to this effect were given in [3] and [16]. Then (5) is exactly the Cardy entropy

$$S_{CFT} = 2\pi \sqrt{\frac{c|\hat{q}_0|}{6}} \quad (6)$$

that the MSW conformal field theory, with central charge

$$c = 6|n_1 n_2 n_3|, \quad (7)$$

assigns to a state where one of the chiral sectors of the theory carries momentum

$$|\hat{q}_0| = |n_p| = |J_1|. \quad (8)$$

The mass of the black ring is however different than the naive mass for the MSW string: it is renormalized as one increases the ratio R_2/R_1 . Still, we can easily understand this effect in an expansion for large R_1 , which corresponds to small λ, μ_i . If we take into account (1) we find

$$M = \frac{\pi R^2}{4G} \left(2\lambda + 2 \sum_i \mu_i + 3\lambda^2 - \sum_i \mu_i^2 + O(\lambda^3, \mu_i^3) \right). \quad (9)$$

We can rewrite this in terms of the unit masses for M5 branes and momentum excitations,

$$m_5 = R \left(\frac{\pi}{4G} \right)^{2/3}, \quad m_p = \frac{1}{R}, \quad (10)$$

so that

$$M = |n_p| m_p + \sum_i |n_i| m_5 - \frac{G((n_p m_p)^2 + \sum_i (n_i m_5)^2)}{\pi R^2} + O(1/R^3). \quad (11)$$

The leading $O(G^0)$ term is the BPS mass of the MSW string. The first $O(G)$ correction, which appears when the string is curved and supersymmetry is broken, is the attractive potential energy due to the self-interaction among the constituents. It shows the characteristic Newtonian fall-off $\propto 1/R^2$ of the five-dimensional potential energy.

Extremal non-BPS charged black rings Ref. [4] presented a larger class of extremal non-supersymmetric black rings, which in addition to the three M5 dipoles possess conserved charges of three kinds of M2 branes, and angular momenta J_1 and J_2 in the S^1 and S^2 of the ring. Besides the parameters R, μ_i, λ , which again must satisfy (1), the solutions contain three boost parameters α_i related to the M2 charges. They are constrained by the condition

$$\frac{C_\lambda}{1+\lambda} s_1 s_2 s_3 = \frac{C_1}{1-\mu_1} s_1 c_2 c_3 + \frac{C_2}{1-\mu_2} c_1 s_2 c_3 + \frac{C_3}{1-\mu_3} c_1 c_2 s_3 \quad (12)$$

which derives from the absence of Dirac-Misner string singularities, which give rise to global causal pathologies. Here we have defined

$$C_\lambda = \epsilon_\lambda \lambda \sqrt{\frac{1+\lambda}{1-\lambda}}, \quad C_i = \epsilon_i \mu_i \sqrt{\frac{1-\mu_i}{1+\mu_i}}, \quad c_i = \cosh \alpha_i, \quad s_i = \sinh \alpha_i, \quad (13)$$

again with $\epsilon_\lambda, \epsilon_i = \pm 1$. Thus we see that the solutions contain six independent continuous parameters. These are *not* enough to allow independent variation of the eight physical parameters of the extremal black rings, namely the M2 and M5 numbers $N_{1,2,3}, n_{1,2,3}$, and

the two spins $J_{1,2}$ (the mass is fixed at extremality, though not necessarily at the BPS bound). In particular the BPS limit with three charges and three dipoles can never be achieved within this family. The more general solution that allows this freedom is still to be constructed.

The M2 and M5 brane numbers are

$$N_i = \left(\frac{2\pi R^3}{G} \right)^{2/3} \frac{s_i c_i}{2} \left[\lambda - \mu_i + \mu_j + \mu_k + 2(\mu_j \mu_k + \lambda \mu_i) + \lambda(\mu_i \mu_j + \mu_i \mu_k - \mu_j \mu_k) + \mu_i \mu_j \mu_k \right], \quad (14)$$

$$n_i = \left(\frac{2\pi R^3}{G} \right)^{1/3} \frac{\sqrt{1-\lambda}}{s_i} \left[\prod_{l=1}^3 \sqrt{1+\mu_l} \right] \left(\frac{C_j}{1-\mu_j} s_j c_k + \frac{C_k}{1-\mu_k} c_j s_k \right). \quad (15)$$

Here (ijk) are cyclic permutations of (123) . The angular momentum along the S^1 of the ring is

$$J_1 = \frac{1}{4} \left(\frac{2\pi R^3}{G} \right) (1-\lambda)^{3/2} \left[\prod_{l=1}^3 (1+\mu_l)^{3/2} \right] \times \left(\frac{C_\lambda}{1-\lambda} c_1 c_2 c_3 - \frac{C_1}{1+\mu_1} c_1 s_2 s_3 - \frac{C_2}{1+\mu_2} s_1 c_2 s_3 - \frac{C_3}{1+\mu_3} s_1 s_2 c_3 \right). \quad (16)$$

The angular momentum J_2 on the S^2 is most simply given in terms of the M2 and M5 charges

$$J_2 = -\frac{1}{2} (n_1 N_1 + n_2 N_2 + n_3 N_3). \quad (17)$$

Finally, the horizon entropy is

$$S_{BH} = \pi \left(\frac{2\pi R^3}{G} \right) (1-\lambda) \lambda^{1/2} \left[\prod_{i=1}^3 (1+\mu_i) \mu_i^{1/2} \right] \times \left| \frac{C_\lambda}{\lambda} c_1 c_2 c_3 + \frac{C_1}{\mu_1} c_1 s_2 s_3 + \frac{C_2}{\mu_2} s_1 c_2 s_3 + \frac{C_3}{\mu_3} s_1 s_2 c_3 \right|. \quad (18)$$

After a very long and tedious calculation, we have found that this horizon entropy admits a simple expression in terms of physical parameters,

$$S_{BH} = 2\pi \sqrt{\left| -J_1 n_1 n_2 n_3 - \frac{(N_1 n_1)^2 + (N_2 n_2)^2 + (N_3 n_3)^2}{4} + \frac{N_1 N_2 n_1 n_2 + N_1 N_3 n_1 n_3 + N_2 N_3 n_2 n_3}{2} \right|}. \quad (19)$$

To reproduce this expression using the Cardy entropy (6) with the central charge (7), now we take level

$$\hat{q}_0 = -J_1 - \frac{(N_1 n_1)^2 + (N_2 n_2)^2 + (N_3 n_3)^2}{4n_1 n_2 n_3} + \frac{1}{2} \left(\frac{N_1 N_2}{n_3} + \frac{N_1 N_3}{n_2} + \frac{N_2 N_3}{n_1} \right). \quad (20)$$

Setting $n_p = -J_1$, the momentum in this level is shifted precisely by the momentum excited by the worldvolume fluxes that yield the M2 charges [12]—this zero-mode momentum does not contribute to the degeneracy of the state. A clear derivation of this result can be found in [17]¹.

Thus the MSW theory naturally accounts for the entropy of these black rings. The complexity of the equations, however, has prevented us from obtaining an expression for the mass that extends (11).

However simple this application of the MSW theory looks, it deepens the puzzles posed by previous application of essentially the same formulas to BPS black rings. In [15] the entropy of the supersymmetric ring was matched to an expression for $\hat{q}_0$ essentially like above, but with an additional zero-point contribution $+n_1 n_2 n_3 / 4$ to the momentum. In [14], however, the entropy of *the same* supersymmetric black ring was instead matched to an expression without this zero-point term, but with $\mathcal{N}_i = N_i - n_j n_k$ instead of N_i . Additionally, one should put $-(J_1 + J_2)$ for the momentum units n_p , instead of $-J_1$. None of these oddities appears in the expression we have found for the extremal non-BPS ring, although it may be worth noting that an alternative form for the oscillator level, which also reproduces the correct entropy, is

$$\hat{q}_0 = -(J_1 + J_2) - \frac{(\mathcal{N}_1 n_1)^2 + (\mathcal{N}_2 n_2)^2 + (\mathcal{N}_3 n_3)^2}{4n_1 n_2 n_3} + \frac{1}{2} \left(\frac{\mathcal{N}_1 \mathcal{N}_2}{n_3} + \frac{\mathcal{N}_1 \mathcal{N}_3}{n_2} + \frac{\mathcal{N}_2 \mathcal{N}_3}{n_1} \right) - \frac{3n_1 n_2 n_3}{4}. \quad (21)$$

According to the proposal in [16], the correct expression for n_p should be the momentum associated to the vector tangent to the S^1 at the horizon. This favors the choice $n_p = -(J_1 + J_2)$ in the BPS case (i.e., like in [14]), and it also yields $n_p = -J_1$ for our extremal non-BPS rings². Still, if the proposal of [14] is correct it remains to understand why the M2 charges N_i should be shifted in the BPS case but not in our rings here. Presumably this has to do with the fact that in our extremal non-BPS rings some modes are not excited, and in particular J_2 vanishes when the N_i are zero. These puzzles would be greatly clarified if we possessed the more general eight-parameter extremal solution where $N_{1,2,3}$, $n_{1,2,3}$, $J_{1,2}$ can be varied independently.

Final remarks. To finish, we comment on the fact that both the left and right moving sectors of the CFT are crucial for understanding the state that corresponds to a black ring, even if only one sector is needed to account for the entropy.

Recall that the puzzles discussed above pertain only to black rings with finite radius. As $R \rightarrow \infty$ the formulas of [14], [15], and of the present paper reduce to the same expression,

¹Ref. [17] also describes a class of non-supersymmetric extremal black rings. In that case extremality is achieved by spinning up the rotation of the S^2 , so it appears to be a different effect than in this paper, although a connection may still exist.

²The claim made in [16] that this identification favors $n_p = -J_1$ for BPS rings is incorrect: the tangent to the S^1 of the near-horizon-BTZ-black hole for these black rings is $\partial_\psi + \partial_\phi$ since the coordinate ϕ on the S^2 is shifted by ψ at the horizon.

which accounts for the entropy of a supersymmetric black string with M2 and M5 charges and momentum. In this limit, only the non-supersymmetric sector of the CFT is excited. This identification of the degrees of freedom responsible for the entropy captures only a part of the state, but not the fact that the ring wraps a contractible circle.

Already for a supersymmetric state a difference exists between the cases where the MSW string wraps a Kaluza-Klein circle and where it wraps a (contractible) ring: the right-moving fermions in the former case are in the R sector, while in the latter case they are in the NS sector. While it would be interesting to study the consequences of this in more detail, a more significant difference must exist between the BPS and extremal non-BPS systems: the latter possess ergoregions and hence are unstable to spontaneous superradiance. In [18] (see also [10, 2]) this effect was explained as follows (in the context of fundamental string states): *both* chiral sectors of the theory are excited. One sector has a large degeneracy of states and gives rise to the statistical entropy of the system. The other sector is also excited (unlike in BPS systems), but in a coherent state that gives the string a circular profile and does not contribute to the degeneracy. Pairs of excitations from each of the two sectors can interact and combine to form an outgoing quantum, which necessarily carries angular momentum away. This accounts for spontaneous superradiance. It seems likely that a similar picture should be present, within the context of the MSW string, for the black rings we have considered in this paper. So it should be interesting to understand the excitations of the system that turn the string into a ring.

Acknowledgements. I am indebted to Harvey Reall for discussions that prompted me to reconsider the microscopic calculation of ref. [3] and the point discussed in footnote 2. This work was supported in part by DURSI 2005 SGR 00082, CICYT FPA 2004-04582-C02-02, and the European Community FP6 program MRTN-CT-2004-005104.

References

- [1] D. N. Page, “Particle Emission Rates From A Black Hole. 2. Massless Particles From A Rotating Hole,” *Phys. Rev. D* **14** (1976) 3260.
- [2] O. J. C. Dias, R. Emparan and A. Maccarrone, “Microscopic Theory of Black Hole Superradiance,” *Phys. Rev. D* **77**, 064018 (2008) [arXiv:0712.0791 [hep-th]].
- [3] R. Emparan, “Rotating circular strings, and infinite non-uniqueness of black rings,” *JHEP* **0403** (2004) 064 [arXiv:hep-th/0402149].
- [4] H. Elvang, R. Emparan and P. Figueras, “Non-supersymmetric black rings as thermally excited supertubes,” *JHEP* **0502** (2005) 031 [arXiv:hep-th/0412130].

- [5] For a review, see R. Emparan and H. S. Reall, “Black rings,” *Class. Quant. Grav.* **23** (2006) R169 [arXiv:hep-th/0608012].
- [6] D. Astefanesei, K. Goldstein, R. P. Jena, A. Sen and S. P. Trivedi, “Rotating attractors,” *JHEP* **0610**, 058 (2006) [arXiv:hep-th/0606244].
- [7] K. Goldstein and R. P. Jena, “One entropy function to rule them all,” *JHEP* **0711**, 049 (2007) [arXiv:hep-th/0701221].
- [8] H. S. Reall, “Counting the microstates of a vacuum black ring,” arXiv:0712.3226 [hep-th].
- [9] R. Emparan and G. T. Horowitz, “Microstates of a neutral black hole in M theory,” *Phys. Rev. Lett.* **97**, 141601 (2006) [arXiv:hep-th/0607023].
- [10] R. Emparan and A. Maccarrone, “Statistical description of rotating Kaluza-Klein black holes,” *Phys. Rev. D* **75** (2007) 084006 [arXiv:hep-th/0701150].
- [11] G. T. Horowitz and M. M. Roberts, “Counting the Microstates of a Kerr Black Hole,” *Phys. Rev. Lett.* **99**, 221601 (2007) [arXiv:0708.1346 [hep-th]].
- [12] J. M. Maldacena, A. Strominger and E. Witten, “Black hole entropy in M-theory,” *JHEP* **9712**, 002 (1997) [arXiv:hep-th/9711053].
- [13] H. Elvang, R. Emparan, D. Mateos and H. S. Reall, “A supersymmetric black ring,” *Phys. Rev. Lett.* **93**, 211302 (2004) [arXiv:hep-th/0407065].
- [14] I. Bena and P. Kraus, “Microscopic description of black rings in AdS/CFT,” *JHEP* **0412**, 070 (2004) [arXiv:hep-th/0408186].
- [15] M. Cyrier, M. Guica, D. Mateos and A. Strominger, “Microscopic entropy of the black ring,” *Phys. Rev. Lett.* **94**, 191601 (2005) [arXiv:hep-th/0411187].
- [16] R. Emparan and D. Mateos, “Oscillator level for black holes and black rings,” *Class. Quant. Grav.* **22** (2005) 3575 [arXiv:hep-th/0506110].
- [17] F. Larsen, “Entropy of thermally excited black rings,” *JHEP* **0510** (2005) 100 [arXiv:hep-th/0505152].
- [18] J. J. Blanco-Pillado, R. Emparan and A. Iglesias, “Fundamental Plasmod Strings and Black Rings,” *JHEP* **0801**, 014 (2008) [arXiv:0712.0611 [hep-th]].